

LETTER TO THE EDITOR

Iran J Allergy Asthma Immunol

March 2008; 7(1): 45-46

Role of *Chlamydia pneumoniae* Infection in Asthma in Northeast of Iran

Ali Ahmadi Torshizi¹, Mohammad Tohidi¹, Davood Attaran¹,
Mehrangiz Khaje Karamadin², and Kiarash Ghazvini²

¹ Department of Respiratory Division, Ghaem Hospital,
Mashhad University of Medical Sciences, Mashhad, Iran

² Department of Microbiology and Virology, Ghaem Hospital,
Mashhad University of Medical Sciences, Mashhad, Iran

Received: 22 May 2007; Received in revised form: 10 November
2007; Accepted: 28 October 2007

ABSTRACT

The role of *Chlamydia pneumoniae* in asthma has drawn much attention in recent years. In this study we assessed the prevalence of *C. pneumoniae* infections in patients with chronic stable and acute exacerbation of asthma and compared it with normal population. Twenty adult patients with chronic stable asthma and 21 patients with acute exacerbations of asthma and 41 matched control subjects were studied for presence of *C. pneumoniae* using cell culture. This study suggests that positive results of *C. pneumoniae* culture are associated with both chronic stable and acute exacerbation of asthma. It could be concluded that *C. pneumoniae* is a risk factor for either development or exacerbation of asthma.

Key words: Asthma; *Chlamydia pneumoniae*; Culture

LETTER

The worldwide high prevalence of asthma and the impact of the disease on quality of life, have led to numerous investigations on the cause of this disease. Recently Chlamydial infection has been suggested to participate in the pathophysiology of asthma.¹⁻³ Considering conflicting data about frequency of *C. pneumoniae* in asthmatic subjects and, regarding that no such study has been done in Middle East, we assessed the prevalence of *C. pneumoniae* infections in well defined patients with chronic stable and acute exacerbation of asthma and compared it with general population.

Corresponding Author: Kiarash Ghazvini, MD;
Microbiology and Virology Research Center, Mashhad University of
Medical Sciences, Ahmad abad Ave, Mashhad 91735, Iran. Tel: (+98
511) 8935 125, Fax: (+98 511) 8012 589,
E-mail: ghazvinik@mums.ac.ir

Culture was used to study the role of *Chlamydia pneumoniae* in asthma among 21 patients whom were admitted as asthma exacerbation and 20 patients with chronic stable asthma in Mashhad University of Medical Sciences, Iran during Feb. 2005 to May 2005. Pair matched healthy control subjects were also chosen for each of these groups. All subjects were examined by the physician and routine questionnaire were completed. Nasopharyngeal epithelial cells, which were obtained with a swab, were used for culture. Specimens were inoculated into cycloheximide-treated monolayers of McCoy cells in tissue culture vessels within 1 h of collection. A positive cell culture was defined when any specimen yielded one or more fluorescent-antibody-stained inclusions in either the primary or the second passage. For increasing reliability, all processes were performed blind. In this study the differences between groups were determined using the Chi square test (SPSS 11.5). For all analyses the level of significance was defined as a P value of 0.05 or less.

Among two asthma groups 47.6% and 35% had positive culture results while about 14.3% and 5% of control subjects had positive results. P-value was statistically significant in asthmatic exacerbation patients (Table 1). The same has been true about the chronic stable asthma group in comparison with its control group (Table 1).

In confirming the cause and effect relationship between *C. pneumoniae* and asthma three conditions should be met: (a) higher frequency of *C. pneumoniae* in asthmatic patients in comparison with control subjects, (b) the presence of *C. pneumoniae* and its quantity, must be related to the severity of disease and (c) successful eradication of *C. pneumoniae*, should be accompanied with clinical improvement. While for *C. pneumoniae* neither of these conditions has been met consistently. In this study we assessed the first condition in both exacerbation and chronic stable asthma. Its also worth mentioning that, we did not consider severity of asthma in evaluating *C. pneumoniae* infection rate, while some authors studied *C. pneumoniae* role in severe asthma,⁴⁻⁶ thus we are not to question or discredit the relationship between *C. pneumoniae* and severe asthma pathogenesis.

While many studies raised question about possible association of *C. pneumoniae* seroprevalence and asthma pathogenesis^{7,8} and even some like Brouard *et al*⁹ discredit any association between *C. pneumoniae* and asthma exacerbation, our study like many other studies found a significant relationship between asthma and this microorganism.^{2,10}

More over Araafa *et al*⁶ found *C. pneumoniae* is related with more severe forms of asthma.

In conclusion, considering significant difference of *C. pneumoniae* infection rates, among asthmatic patients with control subjects and also, regarding high *C. pneumoniae* infection rate in case groups, we deduced that *C. pneumoniae* is a contributing factor in asthma pathogenesis, therefore eradication may be beneficial in asthmatic patients.

ACKNOWLEDGEMENTS

The Office of Research Affairs of Mashhad University of Medical Sciences has provided financial support for this research. The authors would like to thank M. Aalami for his valuable assistance in data collection.

Table 1. The result of tests among cases and their control group.

	Mean age (years)	Sex (M/F)	Number of positive result (percent) *	P-value
Asthma exacerbation	44.42±14	8/13	10 (47.6%)	0.019
Control group for asthma exacerbation	44.42±14	8/13	3 (14.3%)	
Chronic stable asthma	42.00±17	7/13	7 (35%)	0.01
Control group for chronic stable asthma	42.00±17	7/13	1 (5%)	

REFERENCES

1. Emre U, Roblin PM, Gelling M, Dumornay W, Rao M, Hammerschlag MR, et al. The association of Chlamydia pneumoniae infection and reactive airway disease in children. Arch Pediatr Adolesc Med 1994; 148(7):727-32.
2. Hahn DL, McDonald R. Can acute Chlamydia pneumoniae respiratory tract infection initiate chronic asthma? Ann Allergy Asthma Immunol 1998; 81(4):339-44.
3. Hahn DL, Dodge RW, Golubjatnikov R. Association of Chlamydia pneumoniae (strain TWAR) infection with wheezing, asthmatic bronchitis, and adult-onset asthma. JAMA 1991; 266(10):225-30.
4. Von HL, Vasankari T, Liippo K, Wahlstrom E, Puolakkainen M. Chlamydia pneumoniae and severity of asthma. Scand J Infect Dis. 2002; 34(1):22-7.
5. Kroegel C, Rodel J, Mock B. Chlamydia pneumoniae, clarithromycin, and severe asthma. Chest 2001; 120:1035-6.
6. Arafa RM, Matta AM, Bassuny EA. Chlamydia pneumoniae infection in asthmatic patients. Egypt J Immunol 2003; 10(1):103-9.
7. Foschino Barbara MP, Resta O, Aliani M, Guido P, Izzo C, Logroscino C, et al. seroprevalence of chronic chlamydia pneumoniae infection in patients affected by chronic stable asthma. Clin Microbiol Infect 2002; 8(6):358-62.
8. Tuuminen T, Edelstein I, Punin A, Kislova N, Stratchounski L. Use of quantitative and objective enzyme immunoassays to investigate the possible association between chlamydia pneumoniae and Mycoplasma pneumoniae antibodies and asthma. Clin microbiol infect 2004; 10(4):345-8.
9. Brouard J, Freymuth F, Toutain F, Bach N, Vabret A, Gouarin S, Petitjean J, Duhamel JF. Role of viral infections and chlamydia pneumoniae and mycoplasma pneumoniae infections in asthma in infants and young children. Epidemiologic study of 118 children. Arch Pediatr 2002; 9(Suppl 3):365-71.
10. Webley WC, Salva PS, Andrzejewski C, Cirino F, West CA, Tilahun Y, et al. The Bronchial Lavage of Pediatric Patients with asthma contains infectious chlamydia. Am J Respir Crit Care Med 2005; 171(10):1083-8.